Effective Teukolsky Equation on the Brane

Sugumi KANNO
Department of Physics, Kyoto University, Kyoto 606-8501, Japan
Jiro SODA
Department of Physics, Kyoto University, Kyoto 606-8501, Japan

arXiv:gr-qc/0312098v1 22 Dec 2003

Abstract

To estimate Kaluza-Klein (KK) corrections on gravitational waves emitted by perturbed rotating black strings, we give the effective Teukolsky equation on the brane which is separable equation and hence numerically manageable.

1. Introduction

We concentrate on a Randall and Sundrum two-brane model [1]. It would be important to clarify how gravitational waves are generated in the perturbed black string system and how effects of the extra dimensions come into the observed signal of gravitational waves. For this purpose, we derive the effective Teukolsky equation on the brane.

2. Effective Teukolsky Equation

The essential point in deriving the effective Teukolsky equation is to investigate the effects of bulk which is encoded in the Weyl tensor $\delta E_{\mu\nu}$ which satisfies

$$\left(\partial_y^2 - \frac{4}{\ell}\partial_y + \frac{4}{\ell^2}\right)\delta E_{\mu\nu} = -e^{2\frac{y}{\ell}}\hat{\mathcal{L}}_{\mu\nu}{}^{\alpha\beta}\delta E_{\alpha\beta} \equiv -e^{2\frac{y}{\ell}}\hat{\mathcal{L}}\delta E_{\mu\nu} \ , \tag{1}$$

where $\hat{\mathcal{L}}_{\mu\nu}{}^{\alpha\beta}$ stands for the Lichnerowicz operator, $\hat{\mathcal{L}}_{\mu\nu}{}^{\alpha\beta} = \Box\delta^\alpha_\mu\delta^\beta_\nu + 2R_\mu{}^\alpha{}_\nu{}^\beta$. Here, the covariant derivative and the Riemann tensor are constructed from the induced metric $g_{\mu\nu}(x)$. Let $\overset{\oplus}{\phi}(x)$ and $\overset{\ominus}{\phi}(x)$ be the scalar fields on each brane which satisfy $\Box\overset{\oplus}{\phi} = \kappa^2/6\overset{\oplus}{T}$, $\Box\overset{\ominus}{\phi} = \kappa^2/6\overset{\ominus}{T}$, respectively. In the Ricci flat space-time, the identity $(\Box\phi)_{|\mu\nu} = \hat{\mathcal{L}}_{\mu\nu}{}^{\alpha\beta}\left(\phi_{|\alpha\beta}\right)$ holds. Thus, the junction conditions on each brane can be rewritten as

$$e^{2\frac{y}{\ell}}\left[e^{-2\frac{y}{\ell}}\delta E_{\mu\nu}\right]_{,y}\Bigg|_{y=0} = -\hat{\mathcal{L}}\left(\overset{\oplus}{\phi}_{|\mu\nu} - g_{\mu\nu}\Box\overset{\oplus}{\phi}\right) - \frac{\kappa^2}{2}\hat{\mathcal{L}}\overset{\oplus}{T}_{\mu\nu} \equiv \hat{\mathcal{L}}\overset{\oplus}{S}_{\mu\nu} \ , \tag{2}$$

©2004 by Universal Academy Press, Inc.

$$e^{2\frac{y}{\ell}}\left[e^{-2\frac{y}{\ell}}\delta E_{\mu\nu}\right]_{,y}\Bigg|_{y=d} = \hat{\mathcal{L}}\left(\overset{\ominus}{\phi}_{|\alpha\beta} - g_{\mu\nu}\Box\overset{\ominus}{\phi}\right) + \frac{\kappa^2}{2}\hat{\mathcal{L}}\overset{\ominus}{T}_{\mu\nu} \equiv -\hat{\mathcal{L}}\overset{\ominus}{S}_{\mu\nu} \ . \tag{3}$$

The scalar fields $\overset{\oplus}{\phi}$ and $\overset{\ominus}{\phi}$ can be interpreted as the brane fluctuation modes.

The Gregory-Laflamme instability occurs if the curvature length scale of the black hole L is less than the Compton wavelength of KK modes $\sim \ell \exp(d/\ell)$. As we are interested in the stable rotating black string, we assume $\epsilon = (\ell/L)^2 \ll 1$. Under this circumstance, we can use the gradient expansion method where the Weyl tensor $E_{\mu\nu}$ is expanded in the order of ϵ [2]. As a consequence of analysis, we obtain the effective Teukolsky equation in the following form [3]

$$\hat{P}\delta\Psi_4 = \hat{Q}\left(8\pi G T_{nm^*} - \delta E_{nm^*}\Big|_{y=0}\right) + \cdots \ , \tag{4}$$

where $\hat{P}$ and $\hat{Q}$ are the operators and

$$\begin{aligned} \delta E_{\mu\nu}\Big|_{y=0} &= \frac{2}{\ell}\frac{\Omega^2}{1-\Omega^2}\left[\overset{\oplus}{S}_{\mu\nu} + \Omega^2\overset{\ominus}{S}_{\mu\nu}\right] \\ &\quad + \left[\frac{1}{2} + \frac{d}{\ell}\frac{1}{\Omega^2-1}\right]\frac{\ell}{1-\Omega^2}\left[\hat{\mathcal{L}}\overset{\oplus}{S}_{\mu\nu} + \Omega^4\hat{\mathcal{L}}\overset{\ominus}{S}_{\mu\nu}\right] + \frac{\ell}{4}\left[\hat{\mathcal{L}}\overset{\oplus}{S}_{\mu\nu} + \Omega^2\hat{\mathcal{L}}\overset{\ominus}{S}_{\mu\nu}\right] \end{aligned} \tag{5}$$

3. Conclusion

We formulated the perturbative formalism around the Ricci flat two-brane system. In particular, using the gradient expansion method, the effective Teukolsky equation is obtained. This can be used for estimating the Kaluza-Klein corrections on the gravitational waves emitted from the perturbed rotating black string. Our effective Teukolsky equation is completely separable, hence the numerical scheme can be developed in a similar manner as was done in the case of 4-dimensional Teukolsky equation.

Acknowledgments

This work was supported in part by Grant-in-Aid for Scientific Research Fund of the Ministry of Education, Science and Culture of Japan No. 155476 (SK) and No.14540258 (JS).

References

[1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-th/9906064].

[2] S. Kanno and J. Soda, Phys. Rev. D **66**, 083506 (2002) [arXiv:hep-th/0207029]; *ibid.*, 043526 (2002) [arXiv:hep-th/0205188].

[3] S. Kanno and J. Soda, arXiv:gr-qc/0311074.